



## Obesity and the Brain

Implications and Potential Solutions for Children, Middle-Aged Adults, and Older Adults

Auriel A. Willette, Ph.D., M.S.

Assistant Professor of Food Science and Human Nutrition (ISU)
Courtesy Appointments in Psychology (ISU), Biomedical Sciences (ISU),
and Neurology (U. Iowa)

#### **Outline and Take Homes First (!)**


 Iowa is 13<sup>th</sup> in US and tied for 3<sup>rd</sup> in Midwest for adult obesity rates. Obesity is not about blame or shame. The obesity epidemic is very complicated.



• Obesity is consistently related to frontal lobe atrophy in kids to aged adults, which can impact "higher order" thinking.



• Insulin resistance, stemming from obesity, has bad effects on brain health in middle-aged people, "normal" aged people, and aged people across the Alzheimer's disease spectrum.




 To combat obesity and maintain lean muscle: incentivize fruit consumption, less processed meat consumption, and more moderate activity. Changes can improve brain right away.




• Simple things like walking 2 hours/week may help combat Alzheimer's disease.



## Brain Shrinks/Atrophies Over the Human Lifespan



## Obesity, Metabolic Problems, and Brain Health




## **Obesity is NOT a Character Flaw**

- The idea that gluttony and sloth cause obesity, while accepted by virtually everyone, is based on faulty premises and questionable science.
- Meta analyses (i.e., studies summarizing studies) show that excess calorie consumption and lack of activity have small to moderate effects in predicting obesity.
  - ➤ We had similar television, video game, and eating habits (somewhat) back in the 1990's, when obesity rates were MUCH lower¹. Increased computer use has only increased sedentary time per day by 45 minutes on average.
- McAllister et al. (2010) put forward several novel explanations for the obesity epidemic:
  - Changes in gut microflora with certain diets
  - Novel viruses that can cause fat accumulation
  - > Environmental modification of genes that regulate fat storage
  - > Sleep deprivation
  - Hormonal imbalances


## **Obesity, Health Risks, and Brain Health Risks**







#### **Hippocampus and Obesity**





**Children to Young Adults** 

2 of 9 of studies show more atrophy associated with obesity






Middle-Aged to Aged Adults

9 of 19 of studies show more atrophy associated with obesity

## **Prefrontal Cortex and Obesity**





**Children to Young Adults** 

 6 of 9 of studies show more atrophy associated with obesity






Middle-Aged to Aged Adults

12 of 15 of studies show more atrophy associated with obesity

#### **Conclusions**



#### Hippocampus, memory, and aging

- Obesity in literature shows <u>inconsistent</u> associations
- Literature on obesity and memory is very mixed


#### Prefrontal cortex, "higher order" thinking, and aging

- Obesity in literature shows <u>consistent</u> associations
- Literature on obesity and "higher order" thinking is very clear

#### What exactly does the term atrophy mean here?

- Atrophy not due to neural cell death, but fewer cell-to-cell connections
- Weight loss appears to reverse this brain atrophy and problems with thinking

## **Body Mass and Flexible Thinking**




## Alzheimer's Disease and Obesity Related Problems


## Alzheimer's disease (AD) and the Brain


 In AD, unlike normal aging, the brain atrophies due to massive cell death

- During which brain pathology sets in...
  - Amyloid plaques and tau tangles

- And the brain also uses less glucose to power cells
  - Insulin resistance may play a role
- This is a problem, because by 2050
  - Up to 15.6 million Americans will be affected
  - Up to 1.1 trillion/year in Medicare costs







#### **AD Effects on Thinking and Emotion**

- More frequent or severe episodes of forgetfulness
  - Navigating familiar roads
  - Recent conversations
  - Acquaintances, friends, and eventually family



Depression, anxiety, and especially irritability




- Increased mental inflexibility
  - Difficulty adapting to new situations




#### Our "Greatest Hits" Collection of Recent Findings

#### Middle-Aged, At Risk




**Brain Atrophy** 





**Amyloid Plaques** 


#### **Aged, Across AD Spectrum**



Willette et al., 2013, Diabetes Care
Willette et al., 2015 Alzheimer's & Dementia
Willette et al., 2015 Diabetes
Willette et al., 2016 JAMA Neurology
Klinedinst et al., Alzheimer's and Dementia, in review

## What can help improve brain health in lowa?

## **Body Mass and Moderate Physical Activity**




Moderate activity still CRITICAL for maintaining lean muscle mass




More Moderate Physical Activity (Minutes/Week)




## **Body Mass and Intake of Fruits vs. Meats**











## **Potential Therapy for AD**

Walking





Walking for 1 year in people with AD

#### **Acknowledgements**

#### <u>Lab</u>

Brandon Klinedinst Yoojin Lee Kelsey McLimans Allie Plagman Scott Le Army of undergrads



#### **Funding**









# Additional Slides for Reference




FIGURE 2-12Hours per day spent watching television by age group, 1998–2009, and hours per day spent watching television and using computers for youth aged 8-18

NOTES: TotTV = total television content, which includes watching not only standard television programs but also prerecorded shows, DVDs, and television content on other platforms (e.g., Internet, cell phone, iPod); Comp = does not include time spent using a computer for schoolwork.

SOURCES: Rideout et al., 2010; TVB, 2012.